

Secretome of Intestinal bacilli: A natural guard against pathologies

Ilinskaya O., Ulyanova V., Yarullina D., Gataullin I.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017 Ilinskaya, Ulyanova, Yarullina and Gataullin. Current studies of human gut microbiome usually do not consider the special functional role of transient microbiota, although some of its members remain in the host for a long time and produce broad spectrum of biologically active substances. Getting into the gastrointestinal tract (GIT) with food, water and probiotic preparations, two representatives of Bacilli class, genera *Bacillus* and *Lactobacillus*, colonize epithelium blurring the boundaries between resident and transient microbiota. Despite their minor proportion in the microbiome composition, these bacteria can significantly affect both the intestinal microbiota and the entire body thanks to a wide range of secreted compounds. Recently, insufficiency and limitations of pure genome-based analysis of gut microbiota became known. Thus, the need for intense functional studies is evident. This review aims to characterize the *Bacillus* and *Lactobacillus* in GIT, as well as the functional roles of the components released by these members of microbial intestinal community. Complex of their secreted compounds is referred by us as the "bacillary secretome." The composition of the bacillary secretome, its biological effects in GIT and role in counteraction to infectious diseases and oncological pathologies in human organism is the subject of the review.

<http://dx.doi.org/10.3389/fmicb.2017.01666>

Keywords

Bacillus, Gastrointestinal tract, *Lactobacillus*, Metabolites, Secretome

References

- [1] Abriouel, H., Franz, C. M. A. P., Omar, N. B., and Galvez, A. (2011). Diversity and applications of *Bacillus* bacteriocins. *FEMS Microbiol. Rev.* 35, 201-232. doi: 10.1111/j.1574-6976.2010.00244.x
- [2] Adak, S., Aulak, K. S., and Stuehr, D. J. (2002). Direct evidence for nitric oxide production by a nitric-oxide synthase-like protein from *Bacillus subtilis*. *J. Biol. Chem.* 277, 16167-16171. doi: 10.1074/jbc.M201136200
- [3] Adawi, D., Kasravi, F. B., Molin, G., and Jeppsson, B. (1997). Effect of *Lactobacillus* supplementation with and without arginine on liver damage and bacterial translocation in an acute liver injury model in the rat. *Hepatology* 25, 642-647. doi: 10.1002/hep.510250325
- [4] Ahmadova, A., Todorov, S. D., Hadji-Sfaxi, I., Choiset, Y., Rabesona, H., Messaoudi, S., et al. (2013). Antimicrobial and antifungal activities of *Lactobacillus curvatus* strain isolated from homemade Azerbaijani cheese. *Anaerobe* 20, 42-49. doi: 10.1016/j.anaerobe.2013.01.003
- [5] Alakomi, H. L., Skytta, E., Saarela, M., Mattila-Sandholm, T., Latva-Kala, K., and Helander, I. M. (2000). Lactic acid permeabilizes gram-negative bacteria by disrupting the outer membrane. *Appl. Environ. Microbiol.* 66, 2001-2005. doi: 10.1128/AEM.66.5.2001-2005.2000

- [6] Alcántara, C., Blasco, A., Zúñiga, M., and Monedero, V. (2014). Accumulation of polyphosphate in *Lactobacillus* spp. and its involvement in stress resistance. *Appl. Environ. Microbiol.* 80, 1650-1659. doi: 10.1128/AEM.03997-13
- [7] Almand, E. A., Moore, M. D., Outlaw, J., and Jaykus, L. A. (2017). Human norovirus binding to select bacteria representative of the human gut microbiota. *PLoS ONE* 12:e0173124. doi: 10.1371/journal.pone.0173124
- [8] Alou, M. T., Fournier, P. E., and Raoult, D. (2016). "*Bacillus mediterraneensis*," a new bacterial species isolated from human gut microbiota. *New Microbes New Infect.* 12, 86-87. doi: 10.1016/j.nmni.2016.05.006
- [9] Alvarez-Ordóñez, A., Begley, M., Clifford, T., Deasy, T., Considine, K., O'Connor, P., et al. (2014). Investigation of the antimicrobial activity of *Bacillus licheniformis* strains isolated from retail powdered infant milk formulae. *Probiotics Antimicrob. Proteins* 6, 32-40. doi: 10.1007/s12602-013-9151-1
- [10] Alves, N. J., Turner, K. B., Medintz, I. L., and Walper, S. A. (2016). Protecting enzymatic function through directed packaging into bacterial outer membrane vesicles. *Sci. Rep.* 6:24866. doi: 10.1038/srep24866
- [11] Arias, A. A., Ongena, M., Devreese, B., Terrak, M., Joris, B., and Fickers, P. (2013). Characterization of amylolysin, a novel lantibiotic from *Bacillus amyloliquefaciens* GA1. *PLoS ONE* 8:e83037. doi: 10.1371/journal.pone.0083037
- [12] Avila-Calderón, E. D., Araiza-Villanueva, M. G., Cancino-Diaz, J. C., López-Villegas, E. O., Sriranganathan, N., Boyle, S. M., et al. (2015). Roles of bacterial membrane vesicles. *Arch. Microbiol.* 197, 1-10. doi: 10.1007/s00203-014-1042-7
- [13] Ayed, H. B., Maalej, H., Hmidet, N., and Nasri, M. (2015). Isolation and biochemical characterisation of a bacteriocin-like substance produced by *Bacillus amyloliquefaciens* An6. *J. Glob. Antimicrob. Resist.* 3, 255-261. doi: 10.1016/j.jgar.2015.07.001
- [14] Balkus, J. E., Mitchell, C., Agnew, K., Liu, C., Fiedler, T., Cohn, S. E., et al. (2012). Detection of hydrogen peroxide-producing *Lactobacillus* species in the vagina: a comparison of culture and quantitative PCR among HIV-1 seropositive women. *BMC Infect. Dis.* 12:188. doi: 10.1186/1471-2334-12-188
- [15] Barbosa, J., Caetano, T., and Mendo, S. (2015). Class I and class II lanthipeptides produced by *Bacillus* spp. *J. Nat. Prod.* 78, 2850-2866. doi: 10.1021/np500424y
- [16] Barger, S. R., Hoeffler, B. C., Cubillos-Ruiz, A., Russell, W. K., Russell, D. H., and Straight, P. D. (2012). Imaging secondary metabolism of *Streptomyces* sp. Mg1 during cellular lysis and colony degradation of competing *Bacillus subtilis*. *Antonie Van Leeuwenhoek* 102, 435-445. doi: 10.1007/s10482-012-9769-0
- [17] Barrett, E., Ross, R. P., O'Toole, P. W., Fitzgerald, G. F., and Stanton, C. (2012). γ -Aminobutyric acid production by culturable bacteria from the human intestine. *J. Appl. Microbiol.* 113, 411-417. doi: 10.1111/j.1365-2672.2012.05344.x
- [18] Bäuerl, C., Perez-Martinez, G., Yan, F., Polk, D. B., and Monedero, V. (2010). Functional analysis of the p40 and p75 proteins from *Lactobacillus casei* BL23. *J. Mol. Microbiol. Biotechnol.* 19, 231-241. doi: 10.1159/000322233
- [19] Behzadi, E., Hosseini, H. M., and Imani Fooladi, A. A. (2017). The inhibitory impacts of *Lactobacillus rhamnosus* GG-derived extracellular vesicles on the growth of hepatic cancer cells. *Microb. Pathog.* doi: 10.1016/j.micpath.2017.06.016 [Epub ahead of print]
- [20] Bene, K., Varga, Z., Petrov, V. O., Boyko, N., and Rajnavolgyi, E. (2017). Gut microbiota species can provoke both inflammatory and tolerogenic immune responses in human dendritic cells mediated by retinoic acid receptor alpha ligation. *Front. Immunol.* 8:427. doi: 10.3389/fimmu.2017.00427
- [21] Bérdy, J. (2005). Bioactive microbial metabolites. *J. Antibiot.* 58, 1-26. doi: 10.1038/ja.2005.1
- [22] Beric, T., Stankovic, S., Draganic, V., Kojic, M., Lozo, J., and Fira, D. (2014). Novel antilisterial bacteriocin licheniocin 50.2 from *Bacillus licheniformis* VPS50.2 isolated from soil sample. *J. Appl. Microbiol.* 116, 502-510. doi: 10.1111/jam.12393
- [23] Bermudez-Brito, M., Plaza-Díaz, J., Muñoz-Quezada, S., Gómez-Llrente, C., and Gil, A. (2012). Probiotic mechanisms of action. *Ann. Nutr. Metab.* 61, 160-174. doi: 10.1159/000342079
- [24] Bhattarai, Y., Schmidt, B. A., Linden, D. R., Larson, E. D., Grover, M., Beyder, A., et al. (2017). Human derived gut microbiota modulates colonic secretion in mice by regulating 5-HT3 receptor expression via acetate production. *Am. J. Physiol. Gastrointest. Liver Physiol.* 313, G80-G87. doi: 10.1152/ajpgi.00448.2016
- [25] Biller, S. J., Schubotz, F., Roggensack, S. E., Thompson, A. W., Summons, R. E., and Chisholm, S. W. (2014). Bacterial vesicles in marine ecosystems. *Science* 343, 183-186. doi: 10.1126/science.1243457
- [26] Bird, L. E., Ren, J., Zhang, J., Foxwell, N., Hawkins, A. R., Charles, I. G., et al. (2002). Crystal structure of SANOS, a bacterial nitric oxide synthase oxygenase protein from *Staphylococcus aureus*. *Structure* 10, 1687-1696. doi: 10.1016/S0969-2126(02)00911-5
- [27] Boadu, E., Vaskinn, S., Sundkvist, E., Jaeger, R., and Sager, G. (2001). Inhibition by guanosine cyclic monophosphate (cGMP) analogues of uptake of [(3)H]3',5'-cGMP without stimulation of ATPase activity in human erythrocyte inside-out vesicles. *Biochem. Pharmacol.* 62, 425-429. doi: 10.1016/S0006-2952(01)00682-7

- [28] Brown, L., Kessler, A., Cabezas-Sanchez, P., Luque-Garcia, J. L., and Casadevall, A. (2014). Extracellular vesicles produced by the Gram-positive bacterium *Bacillus subtilis* are disrupted by the lipopeptide surfactin. *Mol. Microbiol.* 93, 183-198. doi: 10.1111/mmi.12650
- [29] Brown, L., Wolf, J. M., Prados-Rosales, R., and Casadevall, A. (2015). Through the wall: extracellular vesicles in Gram-positive bacteria, mycobacteria and fungi. *Nat. Rev. Microbiol.* 13, 620-630. doi: 10.1038/nrmicro3480
- [30] Browne, H. P., Forster, S. C., Anonye, B. O., Kumar, N., Neville, B. A., Stares, M. D., et al. (2016). Culturing of 'unculturable' human microbiota reveals novel taxa and extensive sporulation. *Nature* 533, 543-546. doi: 10.1038/nature17645
- [31] Cabrera-Fuentes, H. A., Alba-Alba, C., Aragones, J., Bernhagen, J., Boisvert, W. A., Bøtker, H. E., et al. (2016). Meeting report from the 2nd International Symposium on New Frontiers in Cardiovascular Research. Protecting the cardiovascular system from ischemia: between bench and bedside. *Basic Res. Cardiol.* 111, 7. doi: 10.1007/s00395-015-0527-0
- [32] Cabrera-Fuentes, H. A., Aslam, M., Saffarzadeh, M., Kolpakov, A., Zelenikhin, P., Preissner, K. T., et al. (2013). Internalization of *Bacillus intermedius* ribonuclease (BINASE) induces human alveolar adenocarcinoma cell death. *Toxicon* 69, 219-226. doi: 10.1016/j.toxicon.2013.03.015
- [33] Camilleri, M. (2006). Probiotics and irritable bowel syndrome: rationale, putative mechanisms, and evidence of clinical efficacy. *J. Clin. Gastroenterol.* 40, 264-269. doi: 10.1097/00004836-200603000-00020
- [34] Carding, S., Verbeke, K., Vipond, D. T., Corfe, B. M., and Owen, L. J. (2015). Dysbiosis of the gut microbiota in disease. *Microb. Ecol. Health Dis.* 26:26191. doi: 10.3402/mehd.v26.26191
- [35] Chauhan, A. K., Maheshwari, D. K., and Bajpai, V. K. (2017). Isolation and preliminary characterization of a bacteriocin-producer *Bacillus* strain inhibiting methicillin resistant *Staphylococcus aureus*. *Acta Biol. Hung.* 68, 208-219. doi: 10.1556/018.68.2017.2.8
- [36] Chen, C., Krishnan, V., Macon, K., Manne, K., Narayana, S. V. L., and Schneewind, O. (2013). Secreted proteases control autolysin-mediated biofilm growth of *Staphylococcus aureus*. *J. Biol. Chem.* 288, 29440-29452. doi: 10.1074/jbc.M113.502039
- [37] Chen, X., and Stewart, P. S. (2002). Role of electrostatic interactions in cohesion of bacterial biofilms. *Appl. Microbiol. Biotechnol.* 59, 718-720. doi: 10.1007/s00253-002-1044-2
- [38] Choi, C. H., Kwon, J. G., Kim, S. K., Myung, S. J., Park, K. S., Sohn, C. I., et al. (2015). Efficacy of combination therapy with probiotics and mosapride in patients with IBS without diarrhea: a randomized, double-blind, placebo-controlled, multicenter, Phase II trial. *Neurogastroenterol. Motil.* 27, 705-716. doi: 10.1111/nmo.12544
- [39] Chopra, L., Singh, G., Choudhary, V., and Sahoo, D. K. (2014). Sonorensin: an antimicrobial peptide, belonging to the heterocycloanthracin subfamily of bacteriocins, from a new marine isolate, *Bacillus sonorensis* MT93. *Appl. Environ. Microbiol.* 80, 2981-2990. doi: 10.1128/AEM.04259-13
- [40] Clemente, J. C., Ursell, L. K., Parfrey, L. W., and Knight, R. (2012). The impact of the gut microbiota on human health: an integrative view. *Cell* 148, 1258-1270. doi: 10.1016/j.cell.2012.01.035
- [41] Collins, F. W., O'Connor, P. M., O'Sullivan, O., Rea, M. C., Hill, C., and Ross, R. P. (2016). Formicin-a novel broad-spectrum two-component lantibiotic produced by *Bacillus paralicheniformis* APC 1576. *Microbiology* 162, 1662-1671. doi: 10.1099/mic.0.000340
- [42] De Vuyst, L., and Leroy, F. (2007). Bacteriocins from lactic acid bacteria: production, purification, and food applications. *J. Mol. Microbiol. Biotechnol.* 13, 194-199. doi: 10.1159/000104752
- [43] den Besten, G., Bleeker, A., Gerding, A., van Eunen, K., Havinga, R., van Dijk, T. H., et al. (2015). Short-chain fatty acids protect against high-fat diet-induced obesity via a PPAR γ -dependent switch from lipogenesis to fat oxidation. *Diabetes Metab. Res. Rev.* 64, 2398-2408. doi: 10.2337/db14-1213
- [44] den Besten, G., Lange, K., Havinga, R., van Dijk, T. H., Gerding, A., van Eunen, K., et al. (2013a). Gut-derived short-chain fatty acids are vividly assimilated into host carbohydrates and lipids. *Am. J. Physiol. Gastrointest. Liver Physiol.* 305, G900-G910. doi: 10.1152/ajpgi.00265.2013
- [45] den Besten, G., van Eunen, K., Groen, A. K., Venema, K., Reijngoud, D. J., and Bakker, B. M. (2013b). The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *J. Lipid Res.* 54, 2325-2340. doi: 10.1194/jlr.R036012
- [46] Derrien, M., and van Hylckama Vlieg, J. E. (2015). Fate, activity, and impact of ingested bacteria within the human gut microbiota. *Trends Microbiol.* 23, 354-366. doi: 10.1016/j.tim.2015.03.002
- [47] Dethlefsen, L., Huse, S., Sogin, M. L., and Relman, D. A. (2008). The pervasive effects of an antibiotic on the human gut microbiota, as revealed by deep 16S rRNA sequencing. *PLoS Biol.* 6:e280. doi: 10.1371/journal.pbio.0060280
- [48] Di Luccia, B., D'Apuzzo, E., Varriale, F., Baccigalupi, L., Ricca, E., and Pollice, A. (2016). *Bacillus megaterium* SF185 induces stress pathways and affects the cell cycle distribution of human intestinal epithelial cells. *Benef. Microbes* 7, 609-620. doi: 10.3920/BM2016.0020
- [49] Dobson, A., Cotter, P. D., Ross, R. P., and Hill, C. (2012). Bacteriocin production: a probiotic trait? *Appl. Environ. Microbiol.* 78, 1-6. doi: 10.1128/AEM.05576-11

- [50] Donaldson, G. P., Lee, S. M., and Mazmanian, S. K. (2016). Gut biogeography of the bacterial microbiota. *Nat. Rev. Microbiol.* 14, 20-32. doi: 10.1038/nrmicro3552
- [51] Donia, M. S., Cimerancic, P., Schulze, C. J., Wieland Brown, L. C., Martin, J., Mitreva, M., et al. (2014). A systematic analysis of biosynthetic gene clusters in the human microbiome reveals a common family of antibiotics. *Cell* 158, 1402-1414. doi: 10.1016/j.cell.2014.08.032
- [52] Dubois, J. Y., Kouwen, T. R., Schurich, A. K., Reis, C. R., Ensing, H. T., Trip, E. N., et al. (2009). Immunity to the bacteriocin sublancin 168 is determined by the SunI (YolF) protein of *Bacillus subtilis*. *Antimicrob. Agents Chemother.* 53, 651-661. doi: 10.1128/AAC.01189-08
- [53] Ebel, B., Lemetals, G., Beney, L., Cachon, R., Sokol, H., Langella, P., et al. (2014). Impact of probiotics on risk factors for cardiovascular diseases. A review. *Crit. Rev. Food Sci. Nutr.* 54, 175-189. doi: 10.1080/10408398.2011.579361
- [54] Etienne-Mesmin, L., Chassaing, B., and Gewirtz, A. T. (2017). Tryptophan: a gut microbiota-derived metabolites regulating inflammation. *World J. Gastrointest. Pharmacol. Ther.* 8, 7-9. doi: 10.4292/wjgpt.v8.i1.7
- [55] Fakhry, S., Sorrentini, I., Ricca, E., De Felice, M., and Baccigalupi, L. (2008). Characterization of spore forming *Bacilli* isolated from the human gastrointestinal tract. *J. Appl. Microbiol.* 105, 2178-2186. doi: 10.1111/j.1365-2672.2008.03934.x
- [56] Fekete, A., Givens, D. I., and Lovegrove, J. A. (2015). Casein-derived lactotripeptides reduce systolic and diastolic blood pressure in a meta-analysis of randomised clinical trials. *Nutrients* 7, 659-681. doi: 10.3390/nu7010659
- [57] Fiedler, S., and Heerklotz, H. (2015). Vesicle leakage reflects the target selectivity of antimicrobial lipopeptides from *Bacillus subtilis*. *Biophys. J.* 109, 2079-2089. doi: 10.1016/j.bpj.2015.09.021
- [58] Fleming, D., and Rumbaugh, K. P. (2017). Approaches to dispersing medical biofilms. *Microorganisms* 5:E15. doi: 10.3390/microorganisms5020015
- [59] Fransen, F., Zagato, E., Mazzini, E., Fosso, B., Manzari, C., El Aidy, S., et al. (2015). BALB/c and C57BL/6 mice differ in polyreactive IgA abundance, which impacts the generation of antigen-specific IgA and microbiota diversity. *Immunity* 43, 527-540. doi: 10.1016/j.immuni.2015.08.011
- [60] Fujiya, M., Musch, M. W., Nakagawa, Y., Hu, S., Alverdy, J., Kohgo, Y., et al. (2007). The *Bacillus subtilis* quorum-sensing molecule CSF contributes to intestinal homeostasis via OCTN2, a host cell membrane transporter. *Cell Host Microbe* 1, 299-308. doi: 10.1016/j.chom.2007.05.004
- [61] Gagnière, J., Raisch, J., Veziant, J., Barnich, N., Bonnet, R., Buc, E., et al. (2016). Gut microbiota imbalance and colorectal cancer. *World J. Gastroenterol.* 22, 501-518. doi: 10.3748/wjg.v22.i2.501
- [62] Gao, Z., Guo, B., Gao, R., Zhu, Q., and Qin, H. (2015). Microbiota dysbiosis is associated with colorectal cancer. *Front. Microbiol.* 6:20. doi: 10.3389/fmicb.2015.00020
- [63] Ghelardi, E., Celandroni, F., Salvetti, S., Gueye, S. A., Lupetti, A., and Senesi, S. (2015). Survival and persistence of *Bacillus clausii* in the human gastrointestinal tract following oral administration as spore-based probiotic formulation. *J. Appl. Microbiol.* 119, 552-559. doi: 10.1111/jam.12848
- [64] Ghoshal, U. C., Shukla, R., and Ghoshal, U. (2017). Small intestinal bacterial overgrowth and irritable bowel syndrome: a bridge between functional organic dichotomy. *Gut Liver* 11, 196-208. doi: 10.5009/gnl16126
- [65] Ghouri, Y. A., Richards, D. M., Rahimi, E. F., Krill, J. T., Jelinek, K. A., Dupont, A. W., et al. (2014). Systematic review of randomized controlled trials of probiotics, prebiotics, and synbiotics in inflammatory bowel disease. *Clin. Exp. Gastroenterol.* 7, 473-487. doi: 10.2147/CEG.S27530
- [66] Giraffa, G., Chanishvili, N., and Widyastuti, Y. (2010). Importance of lactobacilli in food and feed biotechnology. *Res. Microbiol.* 161, 480-487. doi: 10.1016/j.resmic.2010.03.001
- [67] Guo, C. J., Chang, F. Y., Wyche, T. P., Backus, K. M., Acker, T. M., Funabashi, M., et al. (2017). Discovery of reactive microbiota-derived metabolites that inhibit host proteases. *Cell* 168, 517.e18-526.e18. doi: 10.1016/j.cell.2016.12.021
- [68] Gusarov, I., Gautier, L., Smolentseva, O., Shamovsky, I., Eremina, S., Mironov, A., et al. (2013). Bacterial nitric oxide extends the lifespan of *C. elegans*. *Cell* 152, 818-830. doi: 10.1016/j.cell.2012.12.043
- [69] Hammes, W. P., and Hertel, C. (2015). "Lactobacillus," in *Bergey's Manual of Systematics of Archaea and Bacteria*, ed. W. B. Whitman (Hoboken, NJ: John Wiley & Sons), 1-76. doi: 10.1002/9781118960608.gbm00604
- [70] Han, X., Mann, E., Gilbert, S., Guan, Y., Steinbrecher, K. A., Montrose, M. H., et al. (2011). Loss of guanylyl cyclase C (GCC) signaling leads to dysfunctional intestinal barrier. *PLoS ONE* 6:e16139. doi: 10.1371/journal.pone.0016139
- [71] Hannig, G., Tchernychev, B., Kurtz, C. B., Bryant, A. P., Currie, M. G., and Silos-Santiago, I. (2014). Guanylate cyclase-C/cGMP: an emerging pathway in the regulation of visceral pain. *Front. Mol. Neurosci.* 7:31. doi: 10.3389/fnmol.2014.00031
- [72] Harmsen, H. J., Raangs, G. C., He, T., Degener, J. E., and Welling, G. W. (2002). Extensive set of 16S rRNA-based probes for detection of bacteria in human feces. *Appl. Environ. Microbiol.* 68, 2982-2990. doi: 10.1128/AEM.68.6.2982-2990.2002

- [73] Haro, C., Rangel-Zuniga, O. A., Alcala-Diaz, J. F., Gomez-Delgado, F., Perez-Martinez, P., Delgado-Lista, J., et al. (2016). Intestinal microbiota is influenced by gender and body mass index. *PLoS ONE* 11:e0154090. doi: 10.1371/journal.pone.0154090
- [74] Hayes, M., Stanton, C., Slattery, H., O'Sullivan, O., Hill, C., Fitzgerald, G. F., et al. (2007). Casein fermentate of *Lactobacillus animalis* DPC6134 contains a range of novel propeptide angiotensin-converting enzyme inhibitors. *Appl. Environ. Microbiol.* 73, 4658-4667. doi: 10.1128/AEM.00096-07
- [75] Hertzberger, R., Arents, J., Dekker, H. L., Pridmore, R. D., Gysler, C., Kleerebezem, M., et al. (2014). H2O2 production in species of the *Lactobacillus acidophilus* group: a central role for a novel NADH-dependent flavin reductase. *Appl. Environ. Microbiol.* 80, 2229-2239. doi: 10.1128/AEM.04272-13
- [76] Holden, J. K., Kang, S., Hollingsworth, S. A., Li, H., Lim, N., Chen, S., et al. (2015). Structure-based design of bacterial nitric oxide synthase inhibitors. *J. Med. Chem.* 58, 994-1004. doi: 10.1021/jm501723p
- [77] Hong, H. A., To, E., Fakhry, S., Baccigalupi, L., Ricca, E., and Cutting, S. M. (2009). Defining the natural habitat of *Bacillus* spore-formers. *Res. Microbiol.* 160, 375-379. doi: 10.1016/j.resmic.2009.06.006
- [78] Huang, X., Lu, Z., and Yang, S. (2006). Antiviral activity of antimicrobial lipopeptide from *Bacillus subtilis* fmbj against pseudorabies virus, porcine parvovirus, newcastle disease virus and infectious bursal disease virus in vitro. *Int. J. Pept. Res. Ther.* 12, 373-377. doi: 10.1007/s10989-006-9041-4
- [79] Hyronimus, B., Le Marrec, C., and Urdaci, M. C. (1998). Coagulin, a bacteriocin-like inhibitory substance produced by *Bacillus coagulans* I4. *J. Appl. Microbiol.* 85, 42-50. doi: 10.1046/j.1365-2672.1998.00466.x
- [80] Iarullina, D. R., and Ilinskaia, O. N. (2007). Genomic determinants of nitric oxide biosynthesis by *Lactobacillus plantarum*: potential opportunities and reality. *Mol. Biol.* 41, 900-907. doi: 10.1134/S0026893307050159
- [81] Iarullina, D. R., Ilinskaia, O. N., Aganov, A. V., Silkin, N. I., and Zverev, D. G. (2006). Alternative pathways of nitric oxide formation in lactobacilli: EPR evidence for nitric oxide synthase activity. *Mikrobiologiya* 75, 731-736. doi: 10.1134/S0026261706060026
- [82] Ilinskaya, O. N., and Shah Mahmud, R. (2014). Ribonucleases as antiviral agents. *Mol. Biol.* 48, 707-717. doi: 10.1134/S0026893314040050
- [83] Ilinskaya, O. N., Singh, I., Dudkina, E., Ulyanova, V., Kayumov, A., and Barreto, G. (2016). Direct inhibition of oncogenic KRAS by *Bacillus pumilus* ribonuclease (binase). *Biochim. Biophys. Acta* 1863, 1559-1567. doi: 10.1016/j.bbamcr.2016.04.005
- [84] Imlay, J. A. (2003). Pathways of oxidative damage. *Annu. Rev. Microbiol.* 57, 395-418. doi: 10.1146/annurev.micro.57.030502.090938
- [85] Jackson, E. K., Cheng, D., Jackson, T. C., Verrier, J. D., and Gillespie, D. G. (2013). Extracellular guanosine regulates extracellular adenosine levels. *Am. J. Physiol. Cell Physiol.* 304, C406-C421. doi: 10.1152/ajpcell.00212.2012
- [86] Jandhyala, S. M., Talukdar, R., Subramanyam, C., Vuyyuru, H., Sasikala, M., and Reddy, D. N. (2015). Role of the normal gut microbiota. *World J. Gastroenterol.* 21, 8787-8803. doi: 10.3748/wjg.v21.i29.8787
- [87] Jung, T. H., Park, J. H., Jeon, W. M., and Han, K. S. (2015). Butyrate modulates bacterial adherence on LS174T human colorectal cells by stimulating mucin secretion and MAPK signaling pathway. *Nutr. Res. Pract.* 9, 343-349. doi: 10.4162/nrp.2015.9.4.343
- [88] Kaškonienė, V., Stankevičius, M., Bimbiraite-Surviliene, K., Naujokaityte, G., Šerniene, L., Mulkyte, K., et al. (2017). Current state of purification, isolation and analysis of bacteriocins produced by lactic acid bacteria. *Appl. Microbiol. Biotechnol.* 101, 1323-1335. doi: 10.1007/s00253-017-8088-9
- [89] Keller, D., Van Dinter, R., Cash, H., Farmer, S., and Venema, K. (2017). *Bacillus coagulans* GBI-30, 6086 increases plant protein digestion in a dynamic, computer-controlled in vitro model of the small intestine (TIM-1). *Benef. Microbes* 8, 491-496. doi: 10.3920/BM2016.0196
- [90] Khochamit, N., Siripornadulsil, S., Sukon, P., and Siripornadulsil, W. (2015). Antibacterial activity and genotypic-phenotypic characteristics of bacteriocin-producing *Bacillus subtilis* KKU213: potential as a probiotic strain. *Microbiol. Res.* 170, 36-50. doi: 10.1016/j.micres.2014.09.004
- [91] Kim, Y., Edwards, N., and Fenselau, C. (2016). Extracellular vesicle proteomes reflect developmental phases of *Bacillus subtilis*. *Clin. Proteomics* 13, 6. doi: 10.1186/s12014-016-9107-z
- [92] Kinkel, T. L., Ramos-Montañez, S., Pando, J. M., Tadeo, D. V., Strom, E. N., Libby, S. J., et al. (2016). An essential role for bacterial nitric oxide synthase in *Staphylococcus aureus* electron transfer and colonization. *Nat. Microbiol.* 2:16224. doi: 10.1038/nmicrobiol.2016.224
- [93] Konishi, H., Fujiya, M., Tanaka, H., Ueno, N., Moriichi, K., Sasajima, J., et al. (2016). Probiotic-derived ferrichrome inhibits colon cancer progression via JNK-mediated apoptosis. *Nat. Commun.* 7:12365. doi: 10.1038/ncomms12365
- [94] Kumar, R., Grover, S., and Batish, V. K. (2012). Bile salt hydrolase (Bsh) activity screening of lactobacilli: in vitro selection of indigenous *Lactobacillus* strains with potential bile salt hydrolysing and cholesterol-lowering ability. *Probiotics Antimicrob. Proteins* 4, 162-172. doi: 10.1007/s12602-012-9101-3

- [95] Lagier, J. C., Hugon, P., Khelaifia, S., Fournier, P. E., La Scola, B., and Raoult, D. (2015). The rebirth of culture in microbiology through the example of culturomics to study human gut microbiota. *Clin. Microbiol. Rev.* 28, 237-264. doi: 10.1128/CMR.00014-14
- [96] Lagier, J. C., Khelaifia, S., Alou, M. T., Ndongo, S., Dione, N., Hugon, P., et al. (2016). Culture of previously uncultured members of the human gut microbiota by culturomics. *Nat. Microbiol.* 1:16203. doi: 10.1038/nmicrobiol.2016.203
- [97] Lahtinen, S. J. (2012). Probiotic viability-does it matter? *Microb. Ecol. Health Dis.* 23, 10-14. doi: 10.3402/mehd.v23i0.18567
- [98] Lamas, B., Richard, M. L., Leducq, V., Pham, H. P., Michel, M. L., Da Costa, G., et al. (2016). CARD9 impacts colitis by altering gut microbiota metabolism of tryptophan into aryl hydrocarbon receptor ligands. *Nat. Med.* 22, 598-605. doi: 10.1038/nm.4102
- [99] Lan, D., Niu, J., Miao, J., Dong, X., Wang, H., Yang, G., et al. (2016). Expression of guanylate cyclase-C, guanylin, and uroguanylin is downregulated proportionally to the ulcerative colitis disease activity index. *Sci. Rep.* 6:25034. doi: 10.1038/srep25034
- [100] LeBlanc, J. G., Chain, F., Martín, R., Bermúdez-Humarán, L. G., Courau, S., and Langella, P. (2017). Beneficial effects on host energy metabolism of short-chain fatty acids and vitamins produced by commensal and probiotic bacteria. *Microb. Cell Fact.* 16, 79. doi: 10.1186/s12934-017-0691-z
- [101] Lee, M. H., Lee, J., Nam, Y. D., Lee, J. S., Seo, M. J., and Yi, S. H. (2016). Characterization of antimicrobial lipopeptides produced by *Bacillus* sp. LM7 isolated from chungkookjang, a Korean traditional fermented soybean food. *Int. J. Food Microbiol.* 221, 12-18. doi: 10.1016/j.ijfoodmicro.2015.12.010
- [102] Li, M., Lee, K., Hsu, M., Nau, G., Mylonakis, E., and Ramratnam, B. (2017). *Lactobacillus*-derived extracellular vesicles enhance host immune responses against vancomycin-resistant enterococci. *BMC Microbiol.* 17:66. doi: 10.1186/s12866-017-0977-7
- [103] Lim, K. B., Balolong, M. P., Kim, S. H., Oh, J. K., Lee, J. Y., and Kang, D. K. (2016). Isolation and characterization of a broad spectrum bacteriocin from *Bacillus amyloliquefaciens* RX7. *Biomed Res. Int.* 2016:8521476. doi: 10.1155/2016/8521476
- [104] Lin, J. E., Snook, A. E., Li, P., Stoecker, B. A., Kim, G. W., Magee, M. S., et al. (2012). GUCY2C opposes systemic genotoxic tumorigenesis by regulating AKT-dependent intestinal barrier integrity. *PLoS ONE* 7:e31686. doi: 10.1371/journal.pone.0031686
- [105] Liu, G., Kong, Y., Fan, Y., Geng, C., Peng, D., and Sun, M. (2017). Whole-genome sequencing of *Bacillus velezensis* LS69, a strain with a broad inhibitory spectrum against pathogenic bacteria. *J. Biotechnol.* 249, 20-24. doi: 10.1016/j.jbiotec.2017.03.018
- [106] Liu, S., and Weiner, H. L. (2016). Control of the gut microbiome by fecal microRNA. *Microb. Cell* 3, 176-177. doi: 10.15698/mic2016.04.492
- [107] Lopetuso, L. R., Scaldaferri, F., Franceschi, F., and Gasbarrini, A. (2016). *Bacillus clausii* and gut homeostasis: state of the art and future perspectives. *Expert Rev. Gastroenterol. Hepatol.* 10, 943-948. doi: 10.1080/17474124.2016.1200465
- [108] Lubber, J. M., and Kostic, A. D. (2017). Gut microbiota: small molecules modulate host cellular functions. *Curr. Biol.* 27, R307-R310. doi: 10.1016/j.cub.2017.03.026
- [109] Maffei, V. J., Kim, S., Blanchard, E., Luo, M., Jazwinski, S. M., Taylor, C. M., et al. (2017). Biological aging and the human gut microbiota. *J. Gerontol. A Biol. Sci. Med. Sci.* doi: 10.1093/gerona/glx042 [Epub ahead of print]
- [110] Magnúsdóttir, S., Ravcheev, D., de Crécy-Lagard, V., and Thiele, I. (2015). Systematic genome assessment of B-vitamin biosynthesis suggests co-operation among gut microbes. *Front. Genet.* 6:148. doi: 10.3389/fgene.2015.00148
- [111] Majeed, M., Nagabhushanam, K., Natarajan, S., Sivakumar, A., Ali, F., Pande, A., et al. (2016). *Bacillus coagulans* MTCC 5856 supplementation in the management of diarrhea predominant Irritable Bowel Syndrome: a double blind randomized placebo controlled pilot clinical study. *Nutr. J.* 15, 21. doi: 10.1186/s12937-016-0140-6
- [112] Marchesi, J. R., Adams, D. H., Fava, F., Hermes, G. D., Hirschfield, G. M., Hold, G., et al. (2016). The gut microbiota and host health: a new clinical frontier. *Gut* 65, 330-339. doi: 10.1136/gutjnl-2015-309990
- [113] Marco, M. L., Heeney, D., Binda, S., Cifelli, C. J., Cotter, P. D., Foligné, B., et al. (2017). Health benefits of fermented foods: microbiota and beyond. *Curr. Opin. Biotechnol.* 44, 94-102. doi: 10.1016/j.copbio.2016.11.010
- [114] Mariat, D., Firmesse, O., Levenez, F., Guimaraes, V., Sokol, H., Doré, J., et al. (2009). The Firmicutes/Bacteroidetes ratio of the human microbiota changes with age. *BMC Microbiol.* 9:123. doi: 10.1186/1471-2180-9-123
- [115] Martí, J. M., Martínez-Martínez, D., Rubio, T., Gracia, C., Peña, M., Latorre, A., et al. (2017). Health and disease imprinted in the time variability of the human microbiome. *mSystems* 2:e00144-16. doi: 10.1128/mSystems.00144-16

- [116] Martín, R., and Suárez, J. E. (2010). Biosynthesis and degradation of H₂O₂ by vaginal lactobacilli. *Appl. Environ. Microbiol.* 76, 400-405. doi: 10.1128/AEM.01631-09
- [117] Martinez, R. C., Wachsmann, M., Torres, N. I., LeBlanc, J. G., Todorov, S. D., and Franco, B. D. (2013). Biochemical, antimicrobial and molecular characterization of a noncytotoxic bacteriocin produced by *Lactobacillus plantarum* ST71KS. *Food Microbiol.* 34, 376-381. doi: 10.1016/j.fm.2013.01.011
- [118] Mashburn, L. M., and Whiteley, M. (2005). Membrane vesicles traffic signals and facilitate group activities in a prokaryote. *Nature* 437, 422-425. doi: 10.1038/nature03925
- [119] Matsuda, K., Tsuji, H., Asahara, T., Matsumoto, K., Takada, T., and Nomoto, K. (2009). Establishment of an analytical system for the human fecal microbiota, based on reverse transcription-quantitative PCR targeting of multicopy rRNA molecules. *Appl. Environ. Microbiol.* 75, 1961-1969. doi: 10.1128/AEM.01843-08
- [120] Matsuoka, K., and Kanai, T. (2015). The gut microbiota and inflammatory bowel disease. *Semin. Immunopathol.* 37, 47-55. doi: 10.1007/s00281-014-0454-4
- [121] McKay, K. A., Kowalec, K., Brinkman, F., Finlay, B. B., Horwitz, M., Manges, A. R., et al. (2017). From bugs to brains: the microbiome in neurological health. *Mult. Scler. Relat. Disord.* 12, 1-3. doi: 10.1016/j.msard.2016.12.007
- [122] Midha, S., Mishra, R., Aziz, M. A., Sharma, M., Mishra, A., Khandelwal, P., et al. (2005). Cloning, expression and characterization of recombinant nitric oxide synthase-like protein from *Bacillus anthracis*. *Biochem. Biophys. Res. Commun.* 336, 346-356. doi: 10.1016/j.bbrc.2005.08.083
- [123] Mitkevich, V. A., Kretova, O. V., Petrushanko, I. Y., Burnysheva, K. M., Sosin, D. V., Simonenko, O. V., et al. (2013). Ribonuclease binase apoptotic signature in leukemic Kasumi-1 cells. *Biochimie* 95, 1344-1349. doi: 10.1016/j.biochi.2013.02.016
- [124] Morita, H., Yoshikawa, H., Sakata, R., Nagata, Y., and Tanaka, H. (1997). Synthesis of nitric oxide from the two equivalent guanidino nitrogens of L-arginine by *Lactobacillus fermentum*. *J. Bacteriol.* 179, 7812-7815. doi: 10.1128/jb.179.24.7812-7815.1997
- [125] Mourembou, G., Ndjoi-Mbiguino, A., Lekana-Douki, J. B., Fournier, P. E., Raoult, D., and Bittar, F. (2016). Discovery of a new species within *Bacillus* genus in a stool sample from Gabon: "*Bacillus massiliogabonensis*" sp. nov. *New Microbes New Infect.* 15, 3-5. doi: 10.1016/j.nmni.2016.09.013
- [126] Müller, C., Ulyanova, V., Ilinskaya, O., Pleschka, S., and Shah Mahmud, R. (2017). A novel antiviral strategy against MERS-CoV and HCoV-229E using binase to target viral genome replication. *BioNanoScience* 7, 294-299. doi: 10.1007/s12668-016-0341-7
- [127] Müller, S., Strack, S. N., Hoeffler, B. C., Straight, P. D., Kearns, D. B., and Kirby, J. R. (2014). Bacillaene and sporulation protect *Bacillus subtilis* from predation by *Myxococcus xanthus*. *Appl. Environ. Microbiol.* 80, 5603-5610. doi: 10.1128/AEM.01621-14
- [128] Nakamura, K., Sakuragi, N., Takakuwa, A., and Ayabe, T. (2016). Paneth cell α -defensins and enteric microbiota in health and disease. *Biosci. Microbiota Food Health* 35, 57-67. doi: 10.12938/bmfh.2015-019
- [129] Natarajan, N., and Pluznick, J. L. (2014). From microbe to man: the role of microbial short chain fatty acid metabolites in host cell biology. *Am. J. Physiol. Cell Physiol.* 307, C979-C985. doi: 10.1152/ajpcell.00228.2014
- [130] Nava, G. M., and Stappenbeck, T. S. (2011). Diversity of the autochthonous colonic microbiota. *Gut Microbes* 2, 99-104. doi: 10.4161/gmic.2.2.15416
- [131] Nguyen, U. T., and Burrows, L. L. (2014). DNase I and proteinase K impair *Listeria monocytogenes* biofilm formation and induce dispersal of pre-existing biofilms. *Int. J. Food Microbiol.* 187, 26-32. doi: 10.1016/j.ijfoodmicro.2014.06.025
- [132] Nithya, V., Muthukumar, S. P., and Halami, P. M. (2012). Safety assessment of *Bacillus licheniformis* Me1 isolated from milk for probiotic application. *Int. J. Toxicol.* 31, 228-237. doi: 10.1177/1091581812443388
- [133] Nøhr, M. K., Egerod, K. L., Christiansen, S. H., Gille, A., Offermanns, S., Schwartz, T. W., et al. (2015). Expression of the short chain fatty acid receptor GPR41/FFAR3 in autonomic and somatic sensory ganglia. *Neuroscience* 290, 126-137. doi: 10.1016/j.neuroscience.2015.01.040
- [134] Nyangale, E. P., Farmer, S., Cash, H. A., Keller, D., Chernoff, D., and Gibson, G. R. (2015). *Bacillus coagulans* GBI-30, 6086 modulates *Faecalibacterium prausnitzii* in older men and women. *J. Nutr.* 145, 1446-1452. doi: 10.3945/jn.114.199802
- [135] Odamaki, T., Kato, K., Sugahara, H., Hashikura, N., Takahashi, S., Xiao, J. Z., et al. (2016). Age-related changes in gut microbiota composition from newborn to centenarian: a cross-sectional study. *BMC Microbiol.* 16:90. doi: 10.1186/s12866-016-0708-5
- [136] Okamoto, K., Fujiya, M., Nata, T., Ueno, N., Inaba, Y., Ishikawa, C., et al. (2012). Competence and sporulation factor derived from *Bacillus subtilis* improves epithelial cell injury in intestinal inflammation via immunomodulation and cytoprotection. *Int. J. Colorectal Dis.* 27, 1039-1046. doi: 10.1007/s00384-012-1416-8
- [137] Oscariz, J. C., and Pisabarro, A. G. (2001). Classification and mode of action of membrane-active bacteriocins produced by gram-positive bacteria. *Int. Microbiol.* 4, 13-19

- [138] Ott, S. J., Musfeldt, M., Wenderoth, D. F., Hampe, J., Brant, O., Fölsch, U. R., et al. (2004). Reduction in diversity of the colonic mucosa associated bacterial microflora in patients with active inflammatory bowel disease. *Gut* 53, 685-693. doi: 10.1136/gut.2003.025403
- [139] Panda, S., El Khader, I., Casellas, F., López Vivancos, J., García Cors, M., Santiago, A., et al. (2014). Short-term effect of antibiotics on human gut microbiota. *PLoS ONE* 9:e95476. doi: 10.1371/journal.pone.0095476
- [140] Patel, A. K., Singhania, R. R., Pandey, A., and Chincholakar, S. B. (2010). Probiotic bile salt hydrolase: current development and perspectives. *Appl. Biochem. Biotechnol.* 162, 166-180. doi: 10.1007/s12010-009-8738-1
- [141] Pattison, A. M., Merlino, D. J., Blomain, E. S., and Waldman, S. A. (2016). Guanylyl cyclase C signaling axis and colon cancer prevention. *World J. Gastroenterol.* 22, 8070-8077. doi: 10.3748/wjg.v22.i36.8070
- [142] Peng, L., Li, Z. R., Green, R. S., Holzman, I. R., and Lin, J. (2009). Butyrate enhances the intestinal barrier by facilitating tight junction assembly via activation of AMP-activated protein kinase in Caco-2 cell monolayers. *J. Nutr.* 139, 1619-1625. doi: 10.3945/jn.109.104638
- [143] Perez, K. J., Viana, J. D., Lopes, F. C., Pereira, J. Q., Dos Santos, D. M., Oliveira, J. S., et al. (2017). *Bacillus* spp. isolated from puba as a source of biosurfactants and antimicrobial lipopeptides. *Front. Microbiol.* 8:61. doi: 10.3389/fmicb.2017.00061
- [144] Pillai, A., and Nelson, R. (2008). Probiotics for treatment of *Clostridium difficile*-associated colitis in adults. *Cochrane Database Syst. Rev.* 23, CD004611. doi: 10.1002/14651858.CD004611.pub2
- [145] Pridmore, R. D., Pittet, A. C., Praplan, F., and Cavadini, C. (2008). Hydrogen peroxide production by *Lactobacillus johnsonii* NCC 533 and its role in anti-Salmonella activity. *FEMS Microbiol. Lett.* 283, 210-215. doi: 10.1111/j.1574-6968.2008.01176.x
- [146] Prior, I. A., Lewis, P. D., and Mattos, C. (2012). A comprehensive survey of Ras mutations in cancer. *Cancer Res.* 72, 2457-2467. doi: 10.1158/0008-5472.CAN-11-2612
- [147] Quagliarello, A., Aloisio, I., Bozzi Cionci, N., Luiselli, D., D'Auria, G., Martinez-Priego, L., et al. (2016). Effect of *Bifidobacterium breve* on the intestinal microbiota of coeliac children on a gluten free diet: a pilot study. *Nutrients* 8:660. doi: 10.3390/nu8100660
- [148] Rafter, J., Bennett, M., Caderni, G., Clune, Y., Hughes, R., Karlsson, P. C., et al. (2007). Dietary synbiotics reduce cancer risk factors in polypectomized and colon cancer patients. *Am. J. Clin. Nutr.* 85, 488-496
- [149] Rajilic-Stojanovic, M., and de Vos, W. M. (2014). The first 1000 cultured species of the human gastrointestinal microbiota. *FEMS Microbiol. Rev.* 38, 996-1047. doi: 10.1111/1574-6976.12075
- [150] Rapozo, D. C., Bernardazzi, C., and de Souza, H. S. (2017). Diet and microbiota in inflammatory bowel disease: the gut in disharmony. *World J. Gastroenterol.* 23, 2124-2140. doi: 10.3748/wjg.v23.i12.2124
- [151] Reuter, G. (2001). The *Lactobacillus* and *Bifidobacterium* microflora of the human intestine: composition and succession. *Curr. Issues Intest. Microbiol.* 2, 43-53
- [152] Rey, M. W., Ramaiya, P., Nelson, B. A., Brody-Karpin, S. D., Zaretsky, E. J., Tang, M., et al. (2004). Complete genome sequence of the industrial bacterium *Bacillus licheniformis* and comparisons with closely related *Bacillus* species. *Genome Biol.* 5:R77. doi: 10.1186/gb-2004-5-10-r77
- [153] Rhee, K. J., Sethupathi, P., Driks, A., Lanning, D. K., and Knight, K. L. (2004). Role of commensal bacteria in development of gut-associated lymphoid tissues and preimmune antibody repertoire. *J. Immunol.* 172, 1118-1124. doi: 10.4049/jimmunol.172.2.1118
- [154] Ríos-Covián, D., Ruas-Madiedo, P., Margolles, A., Gueimonde, M., de Los Reyes-Gavilán, C. G., and Salazar, N. (2016). Intestinal short chain fatty acids and their link with diet and human health. *Front. Microbiol.* 7:185. doi: 10.3389/fmicb.2016.00185
- [155] Rivera, J., Cordero, R. J., Nakouzi, A. S., Frases, S., Nicola, A., and Casadevall, A. (2010). *Bacillus anthracis* produces membrane-derived vesicles containing biologically active toxins. *Proc. Natl. Acad. Sci. U.S.A.* 107, 19002-19007. doi: 10.1073/pnas.1008843107
- [156] Roier, S., Zingl, F. G., Cakar, F., and Schild, S. (2016). Bacterial outer membrane vesicle biogenesis: a new mechanism and its implications. *Microb. Cell* 3, 257-259. doi: 10.15698/mic2016.06.508
- [157] Rowland, I., Gibson, G., Heinken, A., Scott, K., Swann, J., Thiele, I., et al. (2017). Gut microbiota functions: metabolism of nutrients and other food components. *Eur. J. Nutr.* doi: 10.1007/s00394-017-1445-8 [Epub ahead of print]
- [158] Ruiz, L., Hevia, A., Bernardo, D., Margolles, A., and Sanchez, B. (2014). Extracellular molecular effectors mediating probiotic attributes. *FEMS Microbiol. Lett.* 359, 1-11. doi: 10.1111/1574-6968.12576
- [159] Sakatani, A., Fujiya, M., Ueno, N., Kashima, S., Sasajima, J., Moriichi, K., et al. (2016). Polyphosphate derived from *Lactobacillus brevis* inhibits colon cancer progression through induction of cell apoptosis. *Anticancer. Res.* 36, 591-598
- [160] Salard, I., Mercey, E., Rekka, E., Boucher, J.-L., Nioche, P., Mikula, I., et al. (2006). Analogies and surprising differences between recombinant nitric oxide synthase-like proteins from *Staphylococcus aureus* and *Bacillus anthracis* in their interactions with L-arginine analogs and iron ligands. *J. Inorg. Biochem.* 100, 2024-2033. doi: 10.1016/j.jinorgbio.2006.09.017

- [161] Sandrini, S., Aldriwesh, M., Alruways, M., and Freestone, P. (2015). Microbial endocrinology: host-bacteria communication within the gut microbiome. *J. Endocrinol.* 225, R21-R34. doi: 10.1530/JOE-14-0615
- [162] Sankar, S. A., Lagier, J. C., Pontarotti, P., Raoult, D., and Fournier, P. E. (2015). The human gut microbiome, a taxonomic conundrum. *Syst. Appl. Microbiol.* 38, 276-286. doi: 10.1016/j.syapm.2015.03.004
- [163] Satish Kumar, R., Kanmani, P., Yuvaraj, N., Paari, K. A., Pattukumar, V., and Arul, V. (2013). Traditional Indian fermented foods: a rich source of lactic acid bacteria. *Int. J. Food Sci. Nutr.* 64, 415-428. doi: 10.3109/09637486.2012.746288
- [164] Sazawal, S., Hiremath, G., Dhingra, U., Malik, P., Deb, S., and Black, R. E. (2006). Efficacy of probiotics in prevention of acute diarrhea: a meta-analysis of masked, randomized, placebo-controlled trials. *Lancet Infect. Dis.* 6, 374-382. doi: 10.1016/S1473-3099(06)70495-9
- [165] Segawa, S., Fujiya, M., Konishi, H., Ueno, N., Kobayashi, N., Shigyo, T., et al. (2011). Probiotic-derived polyphosphate enhances the epithelial barrier function and maintains intestinal homeostasis through integrin-p38 MAPK pathway. *PLoS ONE* 6:e23278. doi: 10.1371/journal.pone.0023278
- [166] Selvam, R., Maheswari, P., Kavitha, P., Ravichandran, M., Sas, B., and Ramchand, C. N. (2009). Effect of *Bacillus subtilis* PB6, a natural probiotic on colon mucosal inflammation and plasma cytokines levels in inflammatory bowel disease. *Indian J. Biochem. Biophys.* 46, 79-85
- [167] Senghor, B., Seck, E. H., Khelaifia, S., Bassène, H., Sokhna, C., Fournier, P. E., et al. (2017). Description of '*Bacillus dakarensis*' sp. nov., '*Bacillus sinesaloumensis*' sp. nov., '*Gracilibacillus timonensis*' sp. nov., '*Halobacillus massiliensis*' sp. nov., '*Lentibacillus massiliensis*' sp. nov., '*Oceanobacillus senegalensis*' sp. nov., '*Oceanobacillus timonensis*' sp. nov., '*Virgibacillus dakarensis*' sp. nov. and '*Virgibacillus marseillensis*' sp. nov., nine halophilic new species isolated from human stool. *New Microbes New Infect.* 17, 45-51. doi: 10.1016/j.nmni.2017.01.010
- [168] Servin, A. L. (2004). Antagonistic activities of lactobacilli and bifidobacteria against microbial pathogens. *FEMS Microbiol. Rev.* 28, 405-440. doi: 10.1016/j.femsre.2004.01.003
- [169] Shah Mahmud, R., Efimova, M., Mostafa, A., Ulyanova, V., and Ilinskaya, O. (2016). Antiviral activity of bacterial extracellular ribonuclease against single-, double stranded RNA and DNA containing viruses in cell cultures. *BioNanoScience* 6, 561-563. doi: 10.1007/s12668-016-0279-9
- [170] Shah Mahmud, R., and Ilinskaya, O. N. (2013). Antiviral activity of binase against the pandemic influenza A (H1N1) virus. *Acta Nat.* 5, 44-51
- [171] Shah Mahmud, R., Müller, C., Romanova, Y., Mostafa, A., Ulyanova, V., Pleschka, S., et al. (2017). Ribonuclease from *Bacillus* acts as an antiviral agent against negative-and positive-sense single stranded human respiratory RNA viruses. *Biomed Res. Int.* 2017:5279065. doi: 10.1155/2017/5279065
- [172] Sharon, G., Sampson, T. R., Geschwind, D. H., and Mazmanian, S. K. (2016). The central nervous system and the gut microbiome. *Cell* 167, 915-932. doi: 10.1016/j.cell.2016.10.027
- [173] Shi, N., Li, N., Duan, X., and Niu, H. (2017). Interaction between the gut microbiome and mucosal immune system. *Mil. Med. Res.* 4, 14. doi: 10.1186/s40779-017-0122-9
- [174] Shields, R. C., Mokhtar, N., Ford, M., Hall, M. J., Burgess, J. G., ElBadawey, M. R., et al. (2013). Efficacy of a marine bacterial nuclease against biofilm forming microorganisms isolated from chronic rhinosinusitis. *PLoS ONE* 8:e55339. doi: 10.1371/journal.pone.0055339
- [175] Shobharani, P., and Halami, P. M. (2014). Cellular fatty acid profile and H⁺-ATPase activity to assess acid tolerance of *Bacillus* sp. for potential probiotic functional attributes. *Appl. Microbiol. Biotechnol.* 98, 9045-9058. doi: 10.1007/s00253-014-5981-3
- [176] Shobharani, P., Padmaja, R. J., and Halami, P. M. (2015). Diversity in the antibacterial potential of probiotic cultures *Bacillus licheniformis* MCC2514 and *Bacillus licheniformis* MCC2512. *Res. Microbiol.* 166, 546-554. doi: 10.1016/j.resmic.2015.06.003
- [177] Shukla, S. D., Budden, K. F., Neal, R., and Hansbro, P. M. (2017). Microbiome effects on immunity, health and disease in the lung. *Clin. Transl. Immunol.* 6, e133. doi: 10.1038/cti.2017.6
- [178] Singh, R. K., Chang, H. W., Yan, D., Lee, K. M., Ucmak, D., Wong, K., et al. (2017). Influence of diet on the gut microbiome and implications for human health. *J. Transl. Med.* 15, 73. doi: 10.1186/s12967-017-1175-y
- [179] Siraj, Y., Nguen, N., Zelenikhin, P. V., Sokolova, E. V., Gataullin, I. G., and Ilinskaya, O. N. (2015). Dysbiosis of cultivable aerobic microbiota tightly associated with colon cancer epithelial cells. *RJPBCS* 6, 1658-1663
- [180] Smaoui, S., Elleuch, L., Bejar, W., Karray-Rebai, I., Ayadi, I., Jaouadi, B., et al. (2010). Inhibition of fungi and gram-negative bacteria by bacteriocin BacTN635 produced by *Lactobacillus plantarum* sp. TN635. *Appl. Biochem. Biotechnol.* 162, 1132-1146. doi: 10.1007/s12010-009-8821-7
- [181] Sobke, A., Klinger, M., Hermann, B., Sachse, S., Nietzsche, S., Makarewicz, O., et al. (2012). The urinary antibiotic 5-nitro-8-hydroxyquinoline (Nitroxoline) reduces the formation and induces the dispersal of *Pseudomonas aeruginosa* biofilms by chelation of iron and zinc. *Antimicrob. Agents Chemother.* 56, 6021-6025. doi: 10.1128/AAC.01484-12

- [182] Sokurenko, J. V., Zelenikhin, P. V., Ulyanova, V. V., Kolpakov, A. I., Muler, D., and Ilinskaya, O. N. (2015). Identification of 2',3'-cGMP as an intermediate of RNA catalytic cleavage by binase and evaluation of its biological action. *Russ. J. Bioorg. Chem.* 41, 31-36. doi: 10.1134/S1068162015010136
- [183] Sornplang, P., and Piyadeatsoontorn, S. (2016). Probiotic isolates from unconventional sources: a review. *J. Anim. Sci. Technol.* 58, 26. doi: 10.1186/s40781-016-0108-2
- [184] Stern, N. J., Svetoch, E. A., Eruslanov, B. V., Perelygin, V. V., Mitsevich, E. V., Mitsevich, I. P., et al. (2006). Isolation of a *Lactobacillus salivarius* strain and purification of its bacteriocin, which is inhibitory to *Campylobacter jejuni* in the chicken gastrointestinal system. *Antimicrob. Agents Chemother.* 50, 3111-3116. doi: 10.1128/AAC.00259-06
- [185] Stubbendieck, R. M., and Straight, P. D. (2016). Multifaceted interfaces of bacterial competition. *J. Bacteriol.* 198, 2145-2155. doi: 10.1128/JB.00275-16
- [186] Sudhamsu, J., and Crane, B. R. (2006). Structure and reactivity of a thermostable prokaryotic nitric-oxide synthase that forms a long-lived oxy-heme complex. *J. Biol. Chem.* 281, 9623-9632. doi: 10.1074/jbc.M510062200
- [187] Sumi, C. D., Yang, B. W., Yeo, I. C., and Hahm, Y. T. (2015). Antimicrobial peptides of the genus *Bacillus*: a new era for antibiotics. *Can. J. Microbiol.* 61, 93-103. doi: 10.1139/cjm-2014-0613
- [188] Tam, N. K., Uyen, N. Q., Hong, H. A., Duc, L. H., Hoa, T. T., Serra, C. R., et al. (2006). The intestinal life cycle of *Bacillus subtilis* and close relatives. *J. Bacteriol.* 188, 2692-2700. doi: 10.1128/JB.188.7.2692-2700.2006
- [189] Tamang, J. P., Shin, D. H., Jung, S. J., and Chae, S. W. (2016). Functional properties of microorganisms in fermented foods. *Front. Microbiol.* 7:578. doi: 10.3389/fmicb.2016.00578
- [190] Tang, W. H., Kitai, T., and Hazen, S. L. (2017). Gut microbiota in cardiovascular health and disease. *Circ. Res.* 120, 1183-1196. doi: 10.1161/CIRCRESAHA.117.309715
- [191] Thomas, C., and Versalovic, J. (2010). Probiotics-host communication: modulation of signaling pathways in the intestine. *Gut Microbes* 1, 148-163. doi: 10.4161/gmic.1.3.11712
- [192] Travers, M. A., Sow, C., Zirah, S., Deregnaucourt, C., Chaouch, S., Queiroz, R. M., et al. (2016). Deconjugated bile salts produced by extracellular bile-salt hydrolase-like activities from the probiotic *Lactobacillus johnsonii* La1 inhibit *Giardia duodenalis* in vitro growth. *Front. Microbiol.* 7:1453. doi: 10.3389/fmicb.2016.01453
- [193] Tropini, C., Earle, K. A., Huang, K. C., and Sonnenburg, J. L. (2017). The Gut microbiome: connecting spatial organization to function. *Cell Host Microbe* 21, 433-442. doi: 10.1016/j.chom.2017.03.010
- [194] Tsilingiri, K., and Rescigno, M. (2013). Postbiotics: What else? *Benef. Microbes* 4, 101-107. doi: 10.3920/BM2012.0046
- [195] Ulyanova, V., Shah Mahmud, R., Dudkina, E., Vershinina, V., Domann, E., and Ilinskaya, O. (2016). Phylogenetic distribution of extracellular guanyl-preferring ribonucleases renews taxonomic status of two *Bacillus* strains. *J. Gen. Appl. Microbiol.* 62, 181-188. doi: 10.2323/jgam.2016.02.005
- [196] Ulyanova, V., Vershinina, V., and Ilinskaya, O. (2011). Barnase and binase: twins with distinct fates. *FEBS J.* 278, 3633-3643. doi: 10.1111/j.1742-4658.2011.08294.x
- [197] Urgesi, R., Casale, C., Pistelli, R., Rapaccini, G. L., and de Vitis, I. (2014). A randomized double-blind placebo-controlled clinical trial on efficacy and safety of association of simethicone and *Bacillus coagulans* (Colinox®) in patients with irritable bowel syndrome. *Eur. Rev. Med. Pharmacol. Sci.* 18, 1344-1353
- [198] Valko, M., Morris, H., and Cronin, M. T. (2005). Metals, toxicity and oxidative stress. *Curr. Med. Chem.* 12, 1161-1208. doi: 10.2174/0929867053764635
- [199] Valladares, R., Bojilova, L., Potts, A. H., Cameron, E., Gardner, C., Lorca, G., et al. (2013). *Lactobacillus johnsonii* inhibits indoleamine 2,3-dioxygenase and alters tryptophan metabolite levels in BioBreeding rats. *FASEB J.* 27, 1711-1720. doi: 10.1096/fj.12-223339
- [200] Van De Guchte, M., Ehrlich, S. D., and Maguin, E. (2001). Production of growth-inhibiting factors by *Lactobacillus delbrueckii*. *J. Appl. Microbiol.* 91, 147-153. doi: 10.1046/j.1365-2672.2001.01369.x
- [201] van de Wouw, M., Schellekens, H., Dinan, T. G., and Cryan, J. F. (2017). Microbiota-gut-brain axis: modulator of host metabolism and appetite. *J. Nutr.* 147, 727-745. doi: 10.3945/jn.116.240481
- [202] Vanhoutte, T., Huys, G., De Brandt, E., and Swings, J. (2004). Temporal stability analysis of the microbiota in human feces by denaturing gradient gel electrophoresis using universal and group-specific 16S rRNA gene primers. *FEMS Microbiol. Ecol.* 48, 437-446. doi: 10.1016/j.femsec.2004.03.001
- [203] Vargas, G., Rocha, J. D., Oliveira, D. L., Albuquerque, P. C., Frases, S., Santos, S. S., et al. (2015). Compositional and immunobiological analyses of extracellular vesicles released by *Candida albicans*. *Cell. Microbiol.* 17, 389-407. doi: 10.1111/cmi.12374
- [204] Vinoj, G., Vaseeharan, B., Thomas, S., Spiers, A. J., and Shanthy, S. (2014). Quorum-quenching activity of the AHL-lactonase from *Bacillus licheniformis* DAHB1 inhibits *Vibrio* biofilm formation in vitro and reduces shrimp intestinal colonisation and mortality. *Mar. Biotechnol.* 16, 707-715. doi: 10.1007/s10126-014-9585-9

- [205] Voltan, S., Martinez, D., Elli, M., Brun, P., Longo, S., Porzionato, A., et al. (2008). Lactobacillus crispatus M247-derived H₂O₂ acts as a signal transducing molecule activating peroxisome proliferator activated receptor- γ in the intestinal mucosa. *Gastroenterology* 135, 1216-1227. doi: 10.1053/j.gastro.2008.07.007
- [206] Vuong, H. E., Yano, J. M., Fung, T. C., and Hsiao, E. Y. (2017). The microbiome and host behavior. *Annu. Rev. Neurosci.* 40, 21-49. doi: 10.1146/annurev-neuro-072116-031347
- [207] Walter, J. (2008). Ecological role of lactobacilli in the gastrointestinal tract: implications for fundamental and biomedical research. *Appl. Environ. Microbiol.* 74, 4985-4996. doi: 10.1128/AEM.00753-08
- [208] Walter, J., Hertel, C., Tannock, G. W., Lis, C. M., Munro, K., and Hammes, W. P. (2001). Detection of Lactobacillus, Pediococcus, Leuconostoc, and Weissella species in human feces by using group-specific PCR primers and denaturing gradient gel electrophoresis. *Appl. Environ. Microbiol.* 67, 2578-2585. doi: 10.1128/AEM.67.6.2578-2585.2001
- [209] Wang, Q., Zhao, X., Chamu, J., and Shanmugam, K. T. (2011). Isolation, characterization and evolution of a new thermophilic Bacillus licheniformis for lactic acid production in mineral salts medium. *Bioresour. Technol.* 102, 8152-8158. doi: 10.1016/j.biortech.2011.06.003
- [210] Wang, R., Islam, B. N., Bridges, A., Sharman, S. K., Hu, M., Hou, Y., et al. (2017). cGMP signaling increases antioxidant gene expression by activating forkhead box O3A in the colon epithelium. *Am. J. Pathol.* 187, 377-389. doi: 10.1016/j.ajpath.2016.10.016
- [211] Watters, C. M., Burton, T., Kirui, D. K., and Millenbaugh, N. J. (2016). Enzymatic degradation of in vitro Staphylococcus aureus biofilms supplemented with human plasma. *Infect. Drug Resist.* 9, 71-78. doi: 10.2147/IDR.S103101
- [212] Wilks, M., Wiggins, R., Whiley, A., Hennessy, E., Warwick, S., Porter, H., et al. (2004). Identification and H₂O₂ production of vaginal lactobacilli from pregnant women at high risk of preterm birth and relation with outcome. *J. Clin. Microbiol.* 42, 713-717. doi: 10.1128/JCM.42.2.713-717.2004
- [213] Winek, K., Dirnagl, U., and Meisel, A. (2016). The gut microbiome as therapeutic target in central nervous system diseases: implications for stroke. *Neurotherapeutics* 13, 762-774. doi: 10.1007/s13311-016-0475-x
- [214] Yan, F., Cao, H., Cover, T. L., Whitehead, R., Washington, M. K., and Polk, D. B. (2007). Soluble proteins produced by probiotic bacteria regulate intestinal epithelial cell survival and growth. *Gastroenterology* 132, 562-575. doi: 10.1053/j.gastro.2006.11.022
- [215] Yan, F., and Polk, D. B. (2002). Probiotic bacterium prevents cytokine-induced apoptosis in intestinal epithelial cells. *J. Biol. Chem.* 277, 50959-50965. doi: 10.1074/jbc.M207050200
- [216] Yan, J., Herzog, J. W., Tsang, K., Brennan, C. A., Bower, M. A., Garrett, W. S., et al. (2016). Gut microbiota induce IGF-1 and promote bone formation and growth. *Proc. Natl. Acad. Sci. U.S.A.* 113, E7554-E7563. doi: 10.1073/pnas.1607235113
- [217] Yang, M., Jang, I. T., Kim, H. J., and Park, J. K. (2016). Bacillus spp. or Bacillus spp.-derived membrane vesicles induce the intrinsic pathways of apoptosis of human colon cancer cell lines. *J. Bacteriol. Virol.* 46, 84-92. doi: 10.4167/jbv.2016.46.2.84
- [218] Yang, N. J., and Chiu, I. M. (2017). Bacterial signaling to the nervous system through toxins and metabolites. *J. Mol. Biol.* 429, 587-605. doi: 10.1016/j.jmb.2016.12.023
- [219] Yang, Y., and Jobin, C. (2014). Microbial imbalance and intestinal pathologies: connections and contributions. *Dis. Model Mech.* 7, 1131-1142. doi: 10.1242/dmm.016428
- [220] Yarullina, D. R., Beuerlein, K., Silkin, N. I., and Ilinskaya, O. N. (2015). Nitric oxide production in probiotic Lactobacillus plantarum: revision of the origin. *JMRR* 1, 5-14
- [221] Yarullina, D. R., Mikheeva, R. O., Sabirullina, G. I., Zelenikhin, P. V., Ilinskaya, O. N., and Sitdikova, G. F. (2016). Role of nitric oxide produced by lactobacilli in relaxation of intestinal smooth muscles. *Bull. Exp. Biol. Med.* 160, 343-346. doi: 10.1007/s10517-016-3166-z
- [222] Ye, J., Wu, W., Li, Y., and Li, L. (2017). Influences of the gut microbiota on DNA methylation and histone modification. *Dig. Dis. Sci.* 62, 1155-1164. doi: 10.1007/s10620-017-4538-6
- [223] Yunes, R. A., Poluektova, E. U., Dyachkova, M. S., Klimina, K. M., Kovtun, A. S., Averina, O. V., et al. (2016). GABA production and structure of gadB/gadC genes in Lactobacillus and Bifidobacterium strains from human microbiota. *Anaerobe* 42, 197-204. doi: 10.1016/j.anaerobe.2016.10.011
- [224] Zacharofa, M. P., and Lovitt, R. W. (2012). Bacteriocins produced by lactic acid bacteria. A review article. *APCBEE Proc.* 2, 50-56
- [225] Zamfir, M., Callewaert, R., Cornea, P. C., Savu, L., Vatafu, I., and De Vuyst, L. (1999). Purification and characterization of a bacteriocin produced by Lactobacillus acidophilus IBB 801. *J. Appl. Microbiol.* 87, 923-931. doi: 10.1046/j.1365-2672.1999.00950.x
- [226] Zelante, T., Iannitti Rossana, G., Cunha, C., De Luca, A., Giovannini, G., Pieraccini, G., et al. (2013). Tryptophan catabolites from microbiota engage aryl hydrocarbon receptor and balance mucosal reactivity via interleukin-22. *Immunity* 39, 372-385. doi: 10.1016/j.immuni.2013.08.003

- [227] Zhang, H. L., Li, W. S., Xu, D. N., Zheng, W. W., Liu, Y., Chen, J., et al. (2016). Mucosa-reparing and microbiota-balancing therapeutic effect of *Bacillus subtilis* alleviates dextrate sulfate sodium-induced ulcerative colitis in mice. *Exp. Ther. Med.* 12, 2554-2562. doi: 10.3892/etm.2016.3686
- [228] Zhao, X., and Kuipers, O. P. (2016). Identification and classification of known and putative antimicrobial compounds produced by a wide variety of Bacillales species. *BMC Genomics* 17:882. doi: 10.1186/s12864-016-3224-y
- [229] Zhou, S., Zhang, A., Yin, H., and Chu, W. (2016). *Bacillus* sp. QSI-1 modulate quorum sensing signals reduce *Aeromonas hydrophila* level and alter gut microbial community structure in fish. *Front. Cell. Infect. Microbiol.* 6:184. doi: 10.3389/fcimb.2016.00184
- [230] Zhu, Q., Jin, Z., Wu, W., Gao, R., Guo, B., Gao, Z., et al. (2014). Analysis of the intestinal lumen microbiota in an animal model of colorectal cancer. *PLoS ONE* 9:e90849. doi: 10.1371/journal.pone.0090849